



Mahdiyeh Shahi<sup>1</sup>, Hannah Marie Brown<sup>1</sup>, Rong Chen<sup>1</sup>, Diogo Garcia<sup>2</sup>, Erik Middlebrooks<sup>3</sup>, Kaisorn Lee Chaichana<sup>2</sup>, Alfredo Quiñones-Hinojosa<sup>2</sup>, R. Graham Cooks<sup>1</sup>

<sup>1</sup> Chemistry Department, Purdue University, West Lafayette, IN  
<sup>2</sup> Department of Neurological Surgery, Mayo Clinic-Jacksonville, Jacksonville, FL, Mayo Clinic-Jacksonville, Jacksonville, FL, <sup>3</sup> Department of Radiology, Mayo Clinic-Jacksonville, Jacksonville, FL


## Introduction




## Other biomarkers for glioma?



## Conclusions




## Methodology & Results



## Acknowledgments

This study is funded by the National Cancer (R33 CA240181-01A1), Waters Corporation (grant number: 40002775), and Purdue University COVID-19 Research Disruption Fund.



## Publications

- Anal Bioanal Chem. 2021. DOI: 10.1007/s00216-021-03593-0.
- J Appl Lab Med. 2021. DOI: 10.1093/jalm/jfaa233.
- Scientific reports. 2022. DOI: 10.1038/s41598-022-22924-4.